

ONCE-DAILY  
**OZEMPIC**<sup>®</sup>  
semaglutide tablets 4 mg, 9 mg

Welcome to  
Ozempic<sup>®</sup> pill

[Click here](#) to learn more about Ozempic<sup>®</sup> pill



Bottles are not actual size.

You may pay as little as **\$25** for up to a 3-month prescription.<sup>a</sup>

Text **SIGNUP** to 21848.<sup>b</sup> See back cover for more information.

<sup>a</sup>Commercially insured patients only. Eligibility and other restrictions apply.

<sup>b</sup>Message and data rates may apply. Check with your mobile service provider. See Terms and Conditions of Use at [Ozempic.com](https://www.ozempic.com).

Please see Important Safety Information throughout. Click for [Prescribing Information](#) and [Medication Guide](#).

# Getting started on Ozempic®

## Ozempic® pill dosing



### Start

#### 1.5 mg

The starting dose is 1.5 mg for the first 30 days to help your body get used to the medicine.



### Step

#### 4 mg

After your 1.5 mg pills are finished, you will be prescribed the 4 mg dose to help lower blood sugar.



### Escalate to a higher dosage

#### 9 mg

After at least 30 days taking the 4 mg dose, your health care provider may prescribe the 9 mg dose if more blood sugar control is needed.

Pills and bottles are not actual size.

## What are Ozempic® tablets?

Ozempic® (semaglutide) tablets 4 mg or 9 mg are a prescription medicine used:

- along with diet and exercise to improve blood sugar (glucose) in adults with type 2 diabetes
- to reduce the risk of major cardiovascular events such as heart attack, stroke or death in adults with type 2 diabetes who are at high risk for these events

It is not known if Ozempic® tablets are safe and effective for use in children.

Please see Important Safety Information throughout.

- 2 Click for [Prescribing Information](#) and [Medication Guide](#).

ONCE-DAILY  
**OZEMPIC**®  
semaglutide tablets 4mg, 9mg

## Important Safety Information

### What is the most important information I should know about Ozempic® tablets?

Ozempic® tablets may cause serious side effects, including:

- **Possible thyroid tumors, including cancer.** Tell your healthcare provider if you get a lump or swelling in your neck, hoarseness, trouble swallowing, or shortness of breath. These may be symptoms of thyroid cancer. In studies with rodents, Ozempic® tablets and medicines that work like Ozempic® tablets caused thyroid tumors, including thyroid cancer. It is not known if Ozempic® tablets will cause thyroid tumors or a type of thyroid cancer called medullary thyroid carcinoma (MTC) in people

### Do not use Ozempic® tablets if:

- you or any of your family have ever had MTC, or if you have an endocrine system condition called Multiple Endocrine Neoplasia syndrome type 2 (MEN 2)
- you have had a serious allergic reaction to semaglutide or any of the ingredients in Ozempic® tablets

### Before using Ozempic® tablets, tell your healthcare provider if you have any other medical conditions, including if you:

- have or have had problems with your pancreas or kidneys
- have a history of vision problems related to your diabetes
- are scheduled to have surgery or other procedures that use anesthesia or deep sleepiness (deep sedation)
- are pregnant or plan to become pregnant. It is not known if Ozempic® tablets will harm your unborn baby. You should stop using Ozempic® tablets 2 months before you plan to become pregnant
- are breastfeeding or plan to breastfeed. Breastfeeding is not recommended during treatment with Ozempic® tablets

Please see Important Safety Information throughout.

- 3 Click for [Prescribing Information](#) and [Medication Guide](#).

ONCE-DAILY  
**OZEMPIC**®  
semaglutide tablets 4mg, 9mg

## How to take Ozempic® pill

Take only one pill every morning exactly as your health care provider tells you to



### Sip

Take the pill when you wake up, first thing in the morning, on an empty stomach, with a small sip of water (up to 4 ounces).



### Go

Start your morning. Wait at least 30 minutes before eating, drinking, or taking other oral medications.

**If you miss a dose, the missed dose should be skipped, and the next dose should be taken the following day.**

**Additional considerations:** Do not take Ozempic® pill with any other liquids besides water. Swallow whole. Do not split, crush, chew, or dissolve Ozempic® pill.

## Important Safety Information

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Ozempic® tablets may affect the way some medicines work, and some medicines may affect the way Ozempic® tablets work.

Please see Important Safety Information throughout.

- 4 Click for [Prescribing Information](#) and [Medication Guide](#).

ONCE-DAILY  
**OZEMPIC**®  
semaglutide tablets 4mg, 9mg

## How to store Ozempic® pill



- Keep pills at room temperature, between 68 °F and 77 °F
- Store pills in the closed Ozempic® pill bottle in a dry place away from moisture. You may not want to keep Ozempic® pill in the bathroom if it gets too hot and steamy
- Keep the blue cap on Ozempic® pill bottle to keep your pills protected
- Do not use a pill organizer or other container to store Ozempic® pill
- Keep out of reach of children

Bottle shown is for the 9 mg dose. The 1.5 mg and 4 mg bottles are also available. Bottle is not actual size.

## Important Safety Information

### What are the possible side effects of Ozempic® tablets?

Ozempic® tablets may cause serious side effects, including:

- **inflammation of your pancreas (pancreatitis).** Stop using Ozempic® tablets and call your healthcare provider right away if you have severe pain in your stomach area (abdomen) that will not go away, with or without nausea or vomiting. Sometimes you may feel the pain from your abdomen to your back
- **changes in vision.** Tell your healthcare provider if you have changes in vision during treatment with Ozempic® tablets
- **low blood sugar (hypoglycemia).** Your risk for getting low blood sugar may be higher if you use Ozempic® tablets with another medicine that can cause low blood sugar, such as a sulfonylurea or insulin. **Signs and symptoms of low blood sugar may include:** dizziness or lightheadedness, blurred vision, anxiety, irritability or mood changes, sweating, slurred speech, hunger, confusion or drowsiness, shakiness, weakness, headache, fast heartbeat, and feeling jittery

Please see Important Safety Information throughout.

- 5 Click for [Prescribing Information](#) and [Medication Guide](#).

ONCE-DAILY  
**OZEMPIC**®  
semaglutide tablets 4mg, 9mg

## Most common side effects of Ozempic® pill

The most common side effects of Ozempic® pill may include nausea, vomiting, diarrhea, stomach (abdominal) pain, decreased appetite, and constipation. Nausea, vomiting, and diarrhea are most common when first starting Ozempic® pill and when your dose increases.

Talk to your health care provider about any side effect that bothers you or does not go away. These are not all the possible side effects of Ozempic® tablets.

[Click here](#) for additional helpful information on possible Ozempic® side effects

## Important Safety Information

What are the possible side effects of Ozempic® tablets? (cont.)

- **dehydration leading to kidney problems.** Diarrhea, nausea, and vomiting may cause a loss of fluids (dehydration), which may cause kidney problems. It is important for you to drink fluids to help reduce your chance of dehydration. Tell your healthcare provider right away if you have nausea, vomiting, or diarrhea that does not go away
- **severe stomach problems.** Stomach problems, sometimes severe, have been reported in people who use Ozempic® tablets. Tell your healthcare provider if you have stomach problems that are severe or will not go away
- **serious allergic reactions.** Stop using Ozempic® tablets and get medical help right away, if you have any symptoms of a serious allergic reaction, including swelling of your face, lips, tongue, or throat; problems breathing or swallowing; severe rash or itching; fainting or feeling dizzy; or very rapid heartbeat

Please see Important Safety Information throughout.

6 Click for [Prescribing Information](#) and [Medication Guide](#).

ONCE-DAILY  
**OZEMPIC**®  
semaglutide tablets 4mg, 9mg

# Ozempic<sup>®</sup> pill support



## Text2Connect

Get access to Text2Connect—a text-based coaching program that gives you reminders, inspiration, and resources to help you start and stay on Ozempic<sup>®</sup>.

Text **SIGNUP** to **21848** to get started.<sup>a</sup>

<sup>a</sup>Message and data rates may apply. Check with your mobile service provider. See Terms and Conditions of Use at [Ozempic.com](https://www.ozempic.com).



## NovoCare<sup>®</sup> Pharmacy ships Ozempic<sup>®</sup>

Get direct shipping through NovoCare<sup>®</sup> Pharmacy—your reliable source for authentic Ozempic<sup>®</sup>.



## Have questions about Ozempic<sup>®</sup>?

You can call **1-866-696-4090** to get live support.

## Important Safety Information

### What are the possible side effects of Ozempic<sup>®</sup> tablets? (cont.)

- **gallbladder problems.** Gallbladder problems have happened in some people who take Ozempic<sup>®</sup> tablets. Tell your healthcare provider right away if you get symptoms of gallbladder problems, which may include: pain in your upper stomach (abdomen), yellowing of skin or eyes (jaundice), fever, or clay-colored stools
- **food or liquid getting into the lungs during surgery or other procedures that use anesthesia or deep sleepiness (deep sedation).** Ozempic<sup>®</sup> tablets may increase the chance of food getting into your lungs during surgery or other procedures. Tell all your healthcare providers that you are taking Ozempic<sup>®</sup> tablets before you are scheduled to have surgery or other procedures

The most common side effects of Ozempic<sup>®</sup> tablets may include nausea, stomach (abdominal) pain, diarrhea, decreased appetite, vomiting, and constipation.

Please see Important Safety Information throughout.

- 7 Click for [Prescribing Information](#) and [Medication Guide](#).

ONCE-DAILY  
**OZEMPIC<sup>®</sup>**  
semaglutide tablets 4mg, 9mg

# Ozempic<sup>®</sup> pill savings

## Ozempic<sup>®</sup> Savings Offers

Commercially covered  
pay as little as

**\$25**

for up to a  
3-month prescription<sup>a</sup>



OR

Uninsured or self-pay,

**SAVINGS  
OFFERS**

available at  
[OzempicSavings.com<sup>b</sup>](https://www.OzempicSavings.com)

**OZEMPIC<sup>®</sup>**  
semaglutide tablets 4mg, 9mg

Text **SIGNUP** to **21848** or visit [OzempicSavings.com](https://www.OzempicSavings.com) to get started with your savings offer.<sup>c</sup>

<sup>a</sup>Pay as little as \$25, subject to a maximum savings of \$100/month. Eligibility and restrictions apply. Government beneficiaries excluded. One month defined as 1 bottle of 30 tablets of Ozempic<sup>®</sup> pill. Novo Nordisk reserves the right to modify or cancel this program at any time. See full terms at [www.savingscardeligibility.com](https://www.savingscardeligibility.com) and [NovoCarePharmacyTerms.com](https://www.NovoCarePharmacyTerms.com).

<sup>b</sup>Eligibility and restrictions apply. Government beneficiaries excluded. See full terms at [www.savingscardeligibility.com](https://www.savingscardeligibility.com) and [NovoCarePharmacyTerms.com](https://www.NovoCarePharmacyTerms.com).

<sup>c</sup>Message and data rates may apply. Check with your mobile service provider. See Terms and Conditions of Use at [Ozempic.com](https://www.Ozempic.com).

Please see Important Safety Information throughout. Click for [Prescribing Information](#) and [Medication Guide](#).



Only Novo Nordisk manufactures FDA-approved semaglutide medicines, like Ozempic<sup>®</sup>.

NovoCare<sup>®</sup> and Ozempic<sup>®</sup> are registered trademarks of Novo Nordisk A/S.

Novo Nordisk is a registered trademark of Novo Nordisk A/S.

© 2026 Novo Nordisk All rights reserved. US26OZMT00032 April 2026

ONCE-DAILY  
**OZEMPIC<sup>®</sup>**  
semaglutide tablets 4mg, 9mg