


You may pay as little as \$25 for up to a 3-month prescription.^a Text **BEGIN** to 21848.^b
See inside for more information.

^aCommercially insured patients only. Eligibility and other restrictions apply.

^bMessage and data rates may apply.



ONCE-WEEKLY
OZEMPIC[®]
semaglutide injection 0.5mg, 1mg, 2mg

INSIDE THIS BOOKLET:



HOW TO USE YOUR PEN



**FREE TEXT-BASED SAVINGS
AND SUPPORT PROGRAM**



**DETAILS ON HOW TO
SAVE ON OZEMPIC[®]**

More at [Ozempic.com](https://www.ozempic.com)

Please see Important Safety Information throughout,
and enclosed Medication Guide and Prescribing Information
at <https://www.novo-pi.com/ozempic.pdf>.

Getting started on Ozempic®

By adding once-weekly Ozempic® to your type 2 diabetes management plan, you and your health care provider have taken an important step in helping to lower your blood sugar.

Questions about Ozempic®?

- Text **BEGIN to 21848** to sign up for a free text-based savings and support program^a
- Talk to a Diabetes Health Coach by calling **1-866-696-4090 Monday–Friday, 9:00 AM to 6:00 PM ET**

^aMessage and data rates may apply.

What is Ozempic®?

Ozempic® (semaglutide) injection 0.5 mg, 1 mg, or 2 mg is an injectable prescription medicine used:

- along with diet and exercise to improve blood sugar (glucose) in adults with type 2 diabetes.
- to reduce the risk of major cardiovascular events such as heart attack, stroke, or death in adults with type 2 diabetes with known heart disease.

It is not known if Ozempic® can be used in people who have had pancreatitis.

Ozempic® is not for use in people with type 1 diabetes. It is not known if Ozempic® is safe and effective for use in children under 18 years of age.

Please see Important Safety Information throughout, and enclosed Prescribing Information and Medication Guide at <https://www.novo-pi.com/ozempic.pdf>.



Actor portrayal.

Important Safety Information

Do not share your Ozempic® pen with other people, even if the needle has been changed. You may give other people a serious infection, or get a serious infection from them.

What is the most important information I should know about Ozempic®?


Ozempic® may cause serious side effects, including:

- **Possible thyroid tumors, including cancer.** Tell your health care provider if you get a lump or swelling in your neck, hoarseness, trouble swallowing, or shortness of breath. These may be symptoms of thyroid cancer. In studies with rodents, Ozempic® and medicines that work like Ozempic® caused thyroid tumors, including thyroid cancer. It is not known if Ozempic® will cause thyroid tumors or a type of thyroid cancer called medullary thyroid carcinoma (MTC) in people.
- Do not use Ozempic® if you or any of your family have ever had MTC, or if you have an endocrine system condition called Multiple Endocrine Neoplasia syndrome type 2 (MEN 2).

ONCE-WEEKLY
OZEMPIC
semaglutide injection 0.5mg, 1mg, 2mg

How to take Ozempic®

Here is the dosing schedule for the Ozempic® pen that delivers doses of 0.25 mg or 0.5 mg:



The beginning dose is 0.25 mg for the first 4 weeks. This will give your body a chance to get used to the medicine. At Week 5, the dose should be increased to 0.5 mg.

Missed a dose?

Take the missed dose as soon as possible, within 5 days after the missed dose. If more than 5 days have passed, skip the missed dose and take your next dose on the regularly scheduled day.

Ozempic® also comes in pens that deliver doses of 1 mg or doses of 2 mg if your health care provider determines that you need additional blood sugar control.

Please see Important Safety Information throughout, and enclosed Prescribing Information and Medication Guide at <https://www.novo-pi.com/ozempic.pdf>.

1 Take Ozempic® once a week, on the same day, exactly as prescribed by your health care provider.

You can take Ozempic® with or without food.

You may change the day of the week you use Ozempic®, as long as your last dose was given 2 or more days before.

How should I use Ozempic®?

- Ozempic® is injected under the skin (subcutaneously) into your stomach (abdomen), thigh, or upper arm. **Do not** inject Ozempic® into a muscle (intramuscularly) or vein (intravenously)
- **Do not** mix insulin and Ozempic® together in the same injection
- Change (rotate) your injection site with each injection. **Do not** use the same site for each injection
- Talk to your health care provider about how to prevent, recognize, and manage low blood sugar (hypoglycemia), high blood sugar (hyperglycemia), and problems you have because of your diabetes

Important Safety Information

Do not use Ozempic® if:

- you or any of your family have ever had MTC or if you have MEN 2.
- you are allergic to semaglutide or any of the ingredients in Ozempic®. See symptoms of serious allergic reaction in "What are the possible side effects of Ozempic®?"

ONCE-WEEKLY
OZEMPIC
semaglutide injection 0.5mg, 1mg, 2mg

How do I use my Ozempic® pen?

This is a Quick Guide.

Read the Instructions for Use and Important Safety Information before using the pen.



1 Prepare your pen
Check your Ozempic® pen
Read the label to check that your pen contains Ozempic®. Make sure that the Ozempic® medicine in your pen is clear and colorless.



2 Attach a new needle
Tear off the paper tab. Push and turn the needle on until it is tight. Pull off both needle caps.



3 Check the Ozempic® flow with each new pen
Turn the dose selector until the dose counter shows the flow check symbol (•• —). Press and hold in the dose button until the dose counter shows 0. Make sure a drop appears at the needle tip.

Text PEN to 43967 to view a demonstration.^a

^aMessage and data rates may apply.

Please see Important Safety Information throughout, and enclosed Prescribing Information and Medication Guide at <https://www.novo-pi.com/ozempic.pdf>.



4 Give your injection
Select your dose
Turn the dose selector until the dose counter shows the dose you need to inject (0.25 mg or 0.5 mg).



5 Inject your dose
Insert the needle into your skin. Press and hold down the dose button. **After** the dose counter reaches 0, slowly count to 6. Remove the needle from your skin.



6 After your injection
Carefully remove the needle and place it in a sharps container. Put the pen cap back on the pen.

Important Safety Information

Before using Ozempic®, tell your health care provider if you have any other medical conditions, including if you:

- have or have had problems with your pancreas or kidneys.
- have a history of diabetic retinopathy.
- are pregnant or breastfeeding or plan to become pregnant or breastfeed. It is not known if Ozempic® will harm your unborn baby or passes into your breast milk. You should stop using Ozempic® 2 months before you plan to become pregnant.

ONCE-WEEKLY
OZEMPIC®
semaglutide injection 0.5mg, 1mg, 2mg

Common side effects

The most common side effects of Ozempic® may include **nausea, diarrhea, vomiting, stomach (abdominal) pain, and constipation.**

If you experience nausea, here are some general nausea tips that you might find helpful:



Eat bland, low-fat foods, like crackers, toast, and rice



Eat foods that contain water, like soups and gelatin



Avoid fried, greasy, or sweet foods



Avoid lying down after you eat



Go outdoors for fresh air



Eat more slowly



Drink clear or ice-cold drinks

You may report side effects to the FDA at 1-800-FDA-1088. Talk to your health care provider about any side effect that bothers you or does not go away.

Please see **Important Safety Information** throughout, and enclosed **Prescribing Information and Medication Guide** at <https://www.novo-pi.com/ozempic.pdf>.



Actor portrayal.

Important Safety Information

Tell your health care provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, herbal supplements, and other medicines to treat diabetes, including insulin or sulfonylureas.

ONCE-WEEKLY
OZEMPIC
semaglutide injection 0.5mg, 1mg, 2mg

How to store your Ozempic[®] pen

Prior to first use



Refrigerated
36 °F to 46 °F

Until expiration date

After first use



Room temperature
59 °F to 86 °F

OR

Refrigerated
36 °F to 46 °F

56 days

Pen shown delivers doses of 0.25 mg or 0.5 mg.

Text PEN to 43967 for more information on using and storing your pen.^a

^aMessage and data rates may apply.

Important Safety Information

What are the possible side effects of Ozempic[®]?

Ozempic[®] may cause serious side effects, including:

- **inflammation of your pancreas (pancreatitis).** Stop using Ozempic[®] and call your health care provider right away if you have severe pain in your stomach area (abdomen) that will not go away, with or without vomiting. You may feel the pain from your abdomen to your back.
- **changes in vision.** Tell your health care provider if you have changes in vision during treatment with Ozempic[®].

Please see Important Safety Information throughout, and enclosed Prescribing Information and Medication Guide at <https://www.novo-pi.com/ozempic.pdf>.



Actor portrayal.

Important Safety Information

What are the possible side effects of Ozempic[®]? (cont'd)

Ozempic[®] may cause serious side effects, including: (cont'd)

- **low blood sugar (hypoglycemia).** Your risk for getting low blood sugar may be higher if you use Ozempic[®] with another medicine that can cause low blood sugar, such as a sulfonylurea or insulin. **Signs and symptoms of low blood sugar may include:** dizziness or lightheadedness, blurred vision, anxiety, irritability or mood changes, sweating, slurred speech, hunger, confusion or drowsiness, shakiness, weakness, headache, fast heartbeat, and feeling jittery.

ONCE-WEEKLY
OZEMPIC
semaglutide injection 0.5mg, 1mg, 2mg

Support you need, when you need it

Ozempic® provides programs, like the Diabetes Health Coach, as well as a wide range of resources available on [Ozempic.com](https://www.novo-pi.com/ozempic.pdf) to help you on your type 2 diabetes journey.



The Diabetes Health Coach

Get live, one-on-one communication with a Diabetes Health Coach. Your Health Coach can:

- Answer questions about Ozempic® via text, phone call, or email
- Hand-select healthy meal and movement tips to support your needs
- Provide personalized motivation to help you set and reach your diabetes goals

Sign up today or call 1-866-696-4090
Monday through Friday, 9:00 AM to 6:00 PM ET, to get started.



Text2Connect

A text-based coaching program that gives you reminders, inspiration, and resources to help you start and stay on Ozempic®. Text BEGIN to 21848 to get started.^a

^aMessage and data rates may apply.



Actor portrayal.

Important Safety Information

What are the possible side effects of Ozempic®? (cont'd)

Ozempic® may cause serious side effects, including: (cont'd)

- **kidney problems (kidney failure).** In people who have kidney problems, diarrhea, nausea, and vomiting may cause a loss of fluids (dehydration), which may cause kidney problems to get worse. It is important for you to drink fluids to help reduce your chance of dehydration.
- **serious allergic reactions.** Stop using Ozempic® and get medical help right away if you have any symptoms of a serious allergic reaction, including swelling of your face, lips, tongue, or throat; problems breathing or swallowing; severe rash or itching; fainting or feeling dizzy; or very rapid heartbeat.
- **gallbladder problems.** Gallbladder problems have happened in some people who take Ozempic®. Tell your health care provider right away if you get symptoms which may include: pain in your upper stomach (abdomen), fever, yellowing of the skin or eyes (jaundice), or clay-colored stools.

The most common side effects of Ozempic® may include nausea, vomiting, diarrhea, stomach (abdominal) pain, and constipation.

Please see Important Safety Information throughout, and enclosed Prescribing Information and Medication Guide at <https://www.novo-pi.com/ozempic.pdf>.

ONCE-WEEKLY
OZEMPIC
semaglutide injection 0.5mg, 1mg, 2mg

Saving with Ozempic®



Ozempic® Savings Card

To get your offer, text*
BEGIN to 21848
or visit OzempicSavings.com
*Message and data rates apply.

PEEL

PAY AS LITTLE AS \$25 PER PRESCRIPTION

Pay as little as \$25 for a 1-month, 2-month, or 3-month supply for up to 24 months.
Maximum savings of \$150 per 1-month, \$300 per 2-month, or \$450 per 3-month supply. Month defined as 28 days.
Eligibility and other restrictions apply. To receive offer, prescription must be for a 1-, 2-, or 3-month supply for commercially insured patients with coverage.

KIT:
OZEMPIC
semaglutide injection 0.5mg, 1mg, 2mg

 Please see full Prescribing Information, including Medication Guide, or go to Ozempic.com.

To get started with your Savings Card, text **BEGIN to 21848** or visit OzempicSavings.com.^a

With commercial insurance, you may be eligible to pay as little as \$25 when you receive a prescription for either a 1-, 2-, or 3-month supply.^b

^aMessage and data rates may apply. Check with your mobile service provider. See Terms and Conditions of Use at Ozempic.com.

^bApplies to eligible, commercially insured patients with coverage for Ozempic®. Month is defined as 28 days. Maximum savings of \$150 for 1-month, \$300 for 2-month, or \$450 for 3-month supply. For full program details and eligibility requirements, visit OzempicSavings.com.

Please see Important Safety Information throughout, and enclosed Prescribing Information and Medication Guide at <https://www.novo-pi.com/ozempic.pdf>.

Actor portrayal.





Scan this QR code with your phone
or visit StartingOzempic.com for a
digital patient starter experience.

Please see Important Safety Information throughout,
and enclosed Medication Guide and Prescribing
Information at <https://www.novo-pi.com/ozempic.pdf>.

Ozempic® is a registered trademark of Novo Nordisk A/S.
Novo Nordisk is a registered trademark of Novo Nordisk A/S.
© 2023 Novo Nordisk Printed in the U.S.A. US230ZM00329 September 2023



ONCE-WEEKLY
OZEMPIC®
semaglutide injection 0.5mg, 1mg, 2mg